Permutations and Combinations

Fundamental Counting Principle

Fundamental Counting Principle can be used determine the number of possible outcomes when there are two or more characteristics.

Fundamental Counting Principle states that if an event has m possible outcomes and another independent event has n possible outcomes, then there are m * n possible outcomes for the two events together. Fundamental Counting Principle

Lets start with a simple example.

A student is to roll a die and flip a coin. How many possible outcomes will there be?

 1H
 2H
 3H
 4H
 5H
 6H
 6*2 = 12 outcomes

 1T
 2T
 3T
 4T
 5T
 6T
 6*2 = 12 outcomes

12 outcomes

Fundamental Counting Principle

For a college interview, Robert has to choose what to wear from the following: 4 slacks, 3 shirts, 2 shoes and 5 ties. How many possible outfits does he have to choose from?

4*3*2*5 = 120 outfits

A Permutation is an arrangement of items in a particular order.

Notice, ORDER MATTERS!

To find the number of Permutations of n items, we can use the Fundamental Counting Principle or factorial notation.

The number of ways to arrange the letters ABC:

Number of choices for first blank?3 _____Number of choices for second blank?3 2 ____Number of choices for third blank?3 2 1

3*2*1 = 6 3! = 3*2*1 = 6ABC ACB BAC BCA CAB CBA

To find the number of Permutations of n items chosen r at a time, you can use the formula for finding P(n,r) or $_{n}P_{r}$ $_{n}p_{r} = \frac{n!}{(n-r)!}$ where $0 \le r \le n$. $_{5}p_{3} = \frac{5!}{(5-3)!} = \frac{5!}{2!} = 5*4*3 = 60$

Practice:

A combination lock will open when the right choice of three numbers (from 1 to 30, inclusive) is selected. How many different lock combinations are possible assuming no number is repeated?



Practice:

A combination lock will open when the right choice of three numbers (from 1 to 30, inclusive) is selected. How many different lock combinations are possible assuming no number is repeated?

$$_{30}p_3 = \frac{30!}{(30-3)!} = \frac{30!}{27!} = 30*29*28 = 24360$$

Permutations on the Calculator

You can use your calculator to find permutations

- To find the number of permutations of 10 items taken 6 at a time $(_{10}P_6)$:
- Type the total number of items
- Go to the MATH menu and arrow over to PRB
- Choose option 2: nPr
- Type the number of items you want to order

10	MATH NUM CPX <mark>1335</mark> 1:rand 20 nPr	10 nPr 6	10 nPr 6 151200
	3:nCr 4:! 5:randInt(6:randNorm(7:randBin(

Practice:

From a club of 24 members, a President, Vice President, Secretary, Treasurer and Historian are to be elected. In how many ways can the offices be filled?



Practice:

From a club of 24 members, a President, Vice President, Secretary, Treasurer and Historian are to be elected. In how many ways can the offices be filled?

$$_{24}p_5 = \frac{24!}{(24-5)!} = \frac{24!}{19!} =$$

24 * 23 * 22 * 21 * 20 = 5,100,480

A Combination is an arrangement of items in which order does not matter.

ORDER DOES NOT MATTER!

Since the order does not matter in combinations, there are fewer combinations than permutations. The combinations are a "subset" of the permutations.

To find the number of Combinations of n items chosen r at a time, you can use the formula

$${}_{n}C_{r} = \frac{n!}{r!(n-r)!}$$
 where $0 \le r \le n$.

To find the number of Combinations of n items chosen r at a time, you can use the formula n!

 ${}_{n}C_{r} = \frac{n}{r!(n-r)!}$ where $0 \le r \le n$.

$$_{5}C_{3} = \frac{5!}{3!(5-3)!} = \frac{5!}{3!2!} =$$

 $\frac{5*4*3*2*1}{3*2*1*2*1} = \frac{5*4}{2*1} = \frac{20}{2} = 10$

Practice:

To play a particular card game, each player is dealt five cards from a standard deck of 52 cards. How many different hands are possible?





Combinations on the Calculator

You can use your calculator to find combinations

- To find the number of combinations of 10 items taken 6 at a time $({}_{10}C_6)$:
- Type the total number of items
- Go to the MATH menu and arrow over to PRB
- Choose option 3: nCr
- Type the number of items you want to order



Practice:

A student must answer 3 out of 5 essay questions on a test. In how many different ways can the student select the questions?





Practice:

A basketball team consists of two centers, five forwards, and four guards. In how many ways can the coach select a starting line up of one center, two forwards, and two guards?



Practice: A basketball team consists of two centers, five forwards, and four guards. In how many ways can the coach select a starting line up of one center, two forwards, and two guards?



Thus, the number of ways to select the starting line up is 2*10*6 = 120.

Probability with Permutations and Combinations

The 25-member senior class council is selecting officers for president, vice president and secretary. Emily would like to be president, David would like to be vice president, and Jenna would like to be secretary. If the offices are filled at random, beginning with president, what is the probability that they are selected for these offices?

- 1/ (₂₅ P ₃)
- 1/ 13800

Probability with Permutations and Combinations

The 25-member senior class council is selecting members for the prom committee. Stephen, Marcus and Sabrina want would like to be on this committee. If the members are selected at random, what is the probability that all three are selected for this committee?

> 1/ (₂₅ C ₃) 1/ 2300